



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$\therefore x = \frac{a^2 + u^2 - v^2}{2a}, \quad y = \frac{(u^2 - ad)(a - d) + dv^2 + af^2 - aw^2}{2af}.$$

$$\begin{aligned} \text{Also, } u^2 + uv + v^2 &= a^2, \\ u^2 + uw + w^2 &= d^2 + f^2, \\ v^2 + vw + w^2 &= (a - d)^2 + f^2. \end{aligned}$$

$$\therefore u = \frac{ad \pm \frac{1}{2}af\sqrt{3}}{\sqrt{(a^2 + d^2 + f^2 - ad \pm af\sqrt{3})}},$$

$$v = \frac{a^2 - ad \pm \frac{1}{2}af\sqrt{3}}{\sqrt{(a^2 + d^2 + f^2 - ad \pm af\sqrt{3})}},$$

The plus sign is applicable to our solution.

$$w = \frac{d^2 + f^2 - ad \pm \frac{1}{2}af\sqrt{3}}{\sqrt{(a^2 + d^2 + f^2 - ad \pm af\sqrt{3})}},$$

Let $d = \frac{1}{2}a$, and $f = \frac{1}{2}a\sqrt{3}$. Hence, $u = \frac{1}{2}a\sqrt{3} = v = w$. Thus the values of x and y are determined, and equations (7) and (8) are solved.


$$\text{Hence } x = \frac{3ad^2 + 3af^2 + 3a^2d + a^2f\sqrt{3} + 4adf\sqrt{3}}{6(a^2 + d^2 + f^2 - ad + af\sqrt{3})} = \frac{1}{2}a, \text{ when } d = \frac{1}{2}a \text{ and } f = \frac{1}{2}a\sqrt{3}.$$

$$y = \frac{3a^2d\sqrt{3} - 3ad^2\sqrt{3} + af^2\sqrt{3} + 3a^2f}{6(a^2 + d^2 + f^2 - ad + af\sqrt{3})} = \frac{1}{2}a\sqrt{3}, \text{ when } d = \frac{1}{2}a, f = \frac{1}{2}a\sqrt{3}.$$

Also solved by C. N. Schmall.

323. Proposed by W. J. GREENSTREET, Marling School, Stroud, England.

S, S' are the foci of two co-vertical parabolas A and B , the axes of which are at right angles. Draw the circle K on SS' as diameter. K is cut in D and E by a straight line parallel to the axis of A such that S' lies midway between it and that axis. Show that the lines $S'D, S'E$ are parallel to the two tangents to A which are normals to B .



II. Solution by (1) PROPOSER, and (2) R. F. DAVIS, M. A.

I°. The axis of B is the tangent at the common vertex V to A . Hence if PQ be normal at P to B , meeting the axis of B in Q , and if PQ is at the same time a tangent to A , it necessarily follows that SQ is perpendicular to PQ and therefore parallel to TP (the tangent at P to B). Or,

II°. $ST = S'P = S'Q$ (letters as above). Draw $YS'D$ perpendicular to the parallels TP, SQ , and DM perpendicular to the axis of B (Y , of course, being on the tangent at V to B). Then $S'M = VS'$, and D is on the circle, SS' diameter, since $\angle SDS' = 90^\circ$.

III. Solution by the PROPOSER.

The foci S, S' of (A) , $y^2=4ax$, and (B) $x^2=4by$, are, respectively, $(a, 0)$, $(0, b)$, and the circle in question has as its equation $x^2+y^2-ax-by=0$.

Since the common vertex V is $(0, 0)$ and S' is $(0, b)$, the point M in which DE cuts the axis of B is $0, 2b$, and the line DE is $y=2b$.

This cuts the circle where $x^2-ax+2b^2=0\dots(1)$, giving D and E . If the roots of this be x_1 and x_2 , D, E are $(x_1, 2b)$, $(x_2, 2b)$, and $S'D, S'E$ have respectively, for their equations,

$$bx-x_1y+x_1b=0 \text{ and } bx-x_2y+x_2b=0.$$

The tangent at $(at^2, 2at)$ to A is $x-ty+at^2=0$.

The normal at $(2bt', at'^2)$ to B is $x+t'y-2bt'-bt'^3=0$.

These are the same line if $1=\frac{-t}{t'}=\frac{-at^2}{bt'(2+t'^2)}$.

$\therefore t=-t'$, and rejecting the values $t=t'=0$ we have $t^2b-at+2b=0\dots(2)$.

The roots of (1) and (2) are the same, and the property is proved.

Perhaps Prof. Zerr will reconsider the solution he offers.

CALCULUS.

263. Proposed by V. M. SPUNAR, M. S., C. E., East Pittsburg, Pa.

Find a point such that the sum of the squares of its distances from n given points shall be a minimum, and prove that the value so found is $1/n$ th part of the sum of the squares of the mutual distances between the n points, taken two and two.

Solution by G. B. M. ZERR, A. M., Ph. D., Philadelphia, Pa.

This problem is proposed and solved in Williamson's *Differential Calculus*, 9th edition, page 192, Art. 157.

Taking one of the points as origin, and the axes rectangular, let (x, y) be the coordinates of the required point. Let $(a_1, b_1); (a_2, b_2); (a_3, b_3); \dots, (a_{n-1}, b_{n-1})$ be the coordinates of the other $(n-1)$ points. Then

$$x^2+y^2+(x-a_1)^2+(y-b_1)^2+(x-a_2)^2+(y-b_2)^2+\dots \\ + (x-a_{n-1})^2+(y-b_{n-1})^2=u=\text{minimum, or}$$

$$nx^2+ny^2-2(a_1+a_2+\dots+a_{n-1})x-2(b_1+b_2+\dots+b_{n-1})y \\ +a_1^2+a_2^2+\dots+a_{n-1}^2+b_1^2+b_2^2+\dots+b_{n-1}^2=u=\text{minimum.}$$

$$\therefore (nx-a_1-a_2-\dots-a_{n-1})dx+(ny-b_1-b_2-\dots-b_{n-1})dy=0.$$

$$\therefore x=\frac{a_1+a_2+\dots+a_{n-1}}{n}, \quad y=\frac{b_1+b_2+\dots+b_{n-1}}{n}.$$